

Knot topology in QCD

L. P. Zou,^{1,2,3} P. M. Zhang,^{1,4} and D. G. Pak^{1,5}

¹*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

²*School of Nuclear Science and Technology, Lanzhou University, Lanzhou 730000, China*

³*University of Chinese Academy of Sciences, Beijing 100049, China*

⁴*Research Center for Hadron and CSR Physics, Lanzhou University, Lanzhou 730000, China*

⁵*Lab. of Few Nucleon Systems, Institute for Nuclear Physics, Ulugbek, 100214, Uzbekistan*

We consider topological structure of classical vacuum solutions in quantum chromodynamics. Topologically non-equivalent vacuum configurations are classified by non-trivial second and third homotopy groups for coset of the color group $SU(N)$ ($N = 2, 3$) under the action of maximal Abelian stability group. Starting with explicit vacuum knot configurations we study possible exact classical solutions as vacuum excitations. Exact analytic non-static knot solution in a simple CP^1 model in Euclidean space-time has been obtained. We construct an ansatz based on knot and monopole topological vacuum structure for searching new solutions in $SU(2)$ and $SU(3)$ QCD. We show that singular knot-like solutions in QCD in Minkowski space-time can be naturally obtained from knot solitons in integrable CP^1 models. A family of Skyrme type low energy effective theories of QCD admitting exact analytic solutions with non-vanishing Hopf charge is proposed.

PACS numbers: 11.15.-q, 14.20.Dh, 12.38.-t, 12.20.-m

Keywords: Knot solitons, quantum chromodynamics, Skyrme model

I. INTRODUCTION

Topological structure of classical solutions in $SU(N)$ Yang-Mills theory implies numerous physical manifestations in such important phenomena in quantum chromodynamics (QCD) as the chiral symmetry breaking and confinement [1]. The most attractive mechanism of the confinement is based on the Meissner effect in dual color superconductor [2–4] where monopole vacuum condensate is generated dynamically due to quantum corrections [5–7]. Dyons represent alternative topological defects which may play an important role as well as monopoles in description of the confinement at zero and finite temperature [8]. Whereas the instanton and monopole solutions correspond to non-trivial topological Chern-Simons and monopole charges, the topological knot configurations with a non-zero Hopf charge represent another topological objects which become essential in various applications in standard QCD and effective Skyrme type theories of QCD in low energy region [9, 10]. It has been found that knot solitons could be good candidates for description of glueball states which can be treated as excitations over the condensed vacuum [11, 12].

The rich topological structure of QCD as a gauge theory is conditioned by the presence of non-trivial homotopy groups $\pi_k(SU(N)/H)$, where the stability subgroup H determines possible coset spaces with different topological properties. The homotopy group $\pi_3(SU(N)) = Z$ describes topological classes of instanton field configurations corresponding to topological Pontryagin index [13–15]. It is well known that instantons realize tunneling between topologically non-equivalent vacuums and provide dominant contribution to chiral symmetry breaking. Another example of manifestation of non-trivial topology in QCD is provided by the second homotopy group $\pi_2(SU(3)/U(1) \times U(1)) = Z \times Z$ which implies Weyl

symmetric structure of vacuum and singular monopole solutions in $SU(3)$ QCD [16–18].

A nice feature of quantum chromodynamics is that gauge connection (potential) allows natural implementation of the color vector $\hat{n}$ in adjoint representation of $SU(N)$ within the formalism of gauge invariant Abelian decomposition suggested first in [19–22] and developed further in [23–25]. The color vector $\hat{n}$ corresponds to generators of the Cartan algebra of the gauge group $SU(N)$ and gives a suitable tool for description of whole topological structure of the gauge theory. A crucial observation has been made that the classical vacuum in QCD can be explicitly constructed in terms of the color vector $\hat{n}$ [26–28]. *This immediately implies that classical vacuum is strongly degenerated and all topologically non-equivalent vacuums are classified by non-trivial homotopy groups $\pi_{2,3}(SU(N)/H)$.* In particular, in the case of $SU(3)$ QCD it has been shown that classical vacuum possesses a non-trivial Weyl symmetric structure described by the second homotopy group $\pi_2(SU(3)/U(1) \times U(1))$ [28]. It should be stressed, that the color vector $\hat{n}$ represents pure topological degrees of freedom, so that we have still a standard QCD. In low energy region, in effective QCD theories like a generalized Faddeev-Skyrme model [11], the vector $\hat{n}$ becomes dynamical. The knowledge of the classical vacuum structure allows to study vacuum excitations in search of possible finite energy topological solutions and make further steps towards understanding fundamental properties of QCD at quantum level.

In the present paper we consider first the topological structure of the classical vacuum in $SU(N)$, ($N = 2, 3$) QCD and study possible manifestations of topological properties related with the homotopy groups $\pi_{2,3}(SU(3)/U(1) \times U(1))$ and $\pi_{2,3}(SU(2)/U(1))$. Starting with known exact knot solutions from the integrable sector of CP^1 models [29, 30] we will construct a vacuum

with knot topology and obtain new analytic classical solutions in CP^1 model, standard QCD (in Euclidean and Minkowski space-time) and effective Skyrme type theory. The paper is organized as follows. In Section II we describe the general topological structure of the classical vacuum in $SU(2)$ and $SU(3)$ QCD. In Section III we consider a simple CP^1 model which can be treated as a restricted QCD with one field variable $\hat{n}$. Exact non-static knot like solution with a finite Euclidean action has been found. Section IV deals with ansatz for possible topological solutions based on classical vacuum made of $\hat{n}$ with general topology. In Section V we obtain analytic singular knot like solutions in QCD in Minkowski space-time. A family of generalized Skyrme type effective theories admitting exact solutions with non-trivial Hopf numbers is proposed in Section VI.

II. TOPOLOGICAL STRUCTURE OF CLASSICAL VACUUM IN QCD

1. Topology of $SU(2)$ QCD vacuum

It has been shown that knot configurations providing minimums of the energy functional in Faddeev-Skyrme model may correspond to vacuums of QCD in maximal Abelian gauge [26]. Later it has been proved that topologically non-equivalent classical vacuums in pure QCD can be constructed explicitly in terms of a color vector $\hat{n}^a$ ($a = 1, 2, 3$) [27]. By this, the vacuum pure gauge fields $\vec{A}_\mu$ with different Chern-Simons numbers are in one-to-one correspondence with color fields $\hat{n}$ of respective Hopf charges.

One should stress, that the color vector $\hat{n}$ in CP^1 models (as well as in Faddeev-Skyrme theory) represents dynamic field variable whereas in QCD the vector field $\hat{n}$ contains only pure topological degrees of freedom. The most appropriate way how to implement the topological degrees of freedom of $\hat{n}$ into the gauge potential while keeping a standard QCD theory is provided by Cho-Duan-Ge gauge invariant Abelian projection [19–22]

$$\vec{A}_\mu = A_\mu \hat{n} - \frac{1}{g} \hat{n} \times \partial_\mu \hat{n} + \vec{X}_\mu, \quad (1)$$

where A_μ and $\vec{X}_\mu$ are the Abelian and off-diagonal gauge potentials, and $\vec{C}_\mu \equiv -\frac{1}{g} \hat{n} \times \partial_\mu \hat{n}$ is a magnetic potential expressed in terms of $\hat{n}$. For simplicity we put the coupling constant g to be equalled one. The vector $\hat{n}$ has a natural origin in the mathematical structure of the gauge theory, it is defined on the coset G/H where the stability group H is defined by Cartan algebra generators of Lie group G . Notice, since the color vector field $\hat{n}$ represents pure topological degrees of freedom it does not increase the number of physical degrees of freedom in the standard QCD. Notice, that a so-called Cho-Faddeev-Niemi-Shabanov decomposition proposed in [23–25] treats the

color vector $\hat{n}$ as a part of the whole gauge potential. So that, such a decomposition leads to a theory which is different from the standard QCD already at classical level [31].

One can choose an orthonormal frame of three color vector fields $\hat{n}_i^a$, ($i = 1, 2, 3$) which plays a role of metric vielbein on the group manifold $SU(2)$. An interesting analogy can be made between the standard QCD with topological degrees of freedom $\hat{n}_i$ and the topological theory of Lorentz gauge gravity [32] where the vielbein field (tetrad) e_μ^a , ($\mu, a = 0, 1, 2, 3$) exists in a topological phase and obtains dynamical content in low energy effective theory containing the Einstein gravity. In a some sense the standard QCD is in a topological phase where the color vector fields $\hat{n}_i$ describes explicitly the topology of all possible mappings of the base space-time to the group space.

One can verify that the color vector $\hat{n}$ satisfies a covariant constancy condition

$$\begin{aligned} \hat{D}_\mu \hat{n} &\equiv (\partial_\mu + \hat{A}_\mu) \hat{n} = 0, \\ \hat{A}_\mu &= A_\mu \hat{n} + \vec{C}_\mu, \end{aligned} \quad (2)$$

where $\hat{A}_\mu$ is a restricted gauge connection. Let us consider a vector magnetic field strength $\vec{H}_{\mu\nu}$ constructed from the magnetic gauge potential $\vec{C}_\mu$

$$\vec{H}_{\mu\nu} = \partial_\mu \vec{C}_\nu - \partial_\nu \vec{C}_\mu + \vec{C}_\mu \times \vec{C}_\nu. \quad (3)$$

Straightforward calculation shows that the color magnetic field $\vec{H}_{\mu\nu}$ is directed along the color vector $\hat{n}$

$$\vec{H}_{\mu\nu} = H_{\mu\nu} \hat{n}. \quad (4)$$

One can check that the differential 2-form $H = dx^\mu \wedge dx^\nu H_{\mu\nu}$ is closed [19, 20]

$$dH = 0. \quad (5)$$

Due to the Poincare lemma the closed two-form H is locally exact. So, the magnetic field $H_{\mu\nu}$ can be expressed explicitly in terms of a dual Abelian magnetic potential $\tilde{C}_\mu$

$$H_{\mu\nu} = \partial_\mu \tilde{C}_\nu - \partial_\nu \tilde{C}_\mu. \quad (6)$$

The definition of the magnetic field strength $H_{\mu\nu}$ implies the existence of a dual magnetic symmetry $\tilde{U}(1)$ which is an essential ingredient of the dual Meissner mechanism of confinement

$$\delta_{\tilde{U}(1)} \tilde{C}_\mu = -i \partial_\mu \tilde{\alpha}. \quad (7)$$

The gauge invariant Abelian decomposition (1) leads to the following split of the gauge field strength into the Abelian and off-diagonal parts

$$\begin{aligned} \vec{F}_{\mu\nu} &= (F_{\mu\nu} + H_{\mu\nu}) \hat{n} + \hat{D}_\mu \vec{X}_\nu \\ &\quad - \hat{D}_\nu \vec{X}_\mu + g \vec{X}_\mu \times \vec{X}_\nu, \end{aligned} \quad (8)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is an Abelian field strength component, and $\vec{X}_\mu$ can be treated as a color source [19, 20].

An important feature of the gauge invariant Abelian projection is that it implies formulation of a restricted QCD which includes only restricted gauge potential $\vec{A}_\mu$. The restricted QCD possesses full $SU(2)$ gauge invariance if we define a covariant group transformation law for the color vector $\hat{n}$. The Lagrangian of the restricted QCD can be written as follows

$$\mathcal{L}^{restr} = -\frac{1}{4}(F_{\mu\nu} + H_{\mu\nu})^2. \quad (9)$$

The Lagrangian has a manifest covariant Abelian structure within the full $SU(2)$ gauge theory. In the absence of the "electric" part, $F_{\mu\nu} = 0$, the Lagrangian reduces to the Lagrangian of a simple CP^1 model.

One can impose covariant constancy conditions on all three basis vector fields $\hat{n}_i$

$$\vec{D}_\mu \hat{n}_i = 0. \quad (10)$$

Acting on the left hand side of the last equation with a covariant derivative $\vec{D}_\nu$ and taking antisymmetrization over indices μ, ν one obtains the integrability condition

$$\vec{F}_{\mu\nu} = 0 \quad (11)$$

which is exactly an equation defining classical vacuum of QCD. An explicit construction of the classical vacuum of QCD in terms of knot configurations of $\hat{n}$ has been found first in [27]

$$\vec{A}_\mu^{vac} = -\vec{C}_\mu \hat{n} + \vec{C}_\mu. \quad (12)$$

This relation establishes connection between a pure gauge potential and color vector field $\hat{n}$ and implies that the classical vacuum configurations can be described by topologically non-equivalent classes of the color field $\hat{n}$. Namely, the topological classes of $\hat{n}$ are determined by two homotopy groups, $\pi_2(SU(2)/U(1))$ and $\pi_3(SU(2)/U(1)) = \pi_3(S^2)$. The first one describes monopole configurations, whereas the second homotopy describes Hopf mapping $\hat{n} : S^3 \rightarrow S^2$ (we assume that the space R^3 is compactified to a three dimensional sphere S^3). So that, all topological non-equivalent classical vacuums are classified by Hopf, Q_H , and monopole, g_m , charges

$$\begin{aligned} Q_H &= \frac{1}{32\pi^2} \int d^3x \epsilon^{ijk} \vec{C}_i H_{jk}, \\ g_m &= \int_{S^2} \vec{H}_{ij} \cdot \hat{n} d\sigma^{ij}. \end{aligned} \quad (13)$$

One can show that the Hopf number equals to the Chern-Simons number for vacuum gauge field configurations A_μ^{vac} constructed from $\hat{n}$.

To study possible physical manifestations of the vacuum structure we will consider explicit expressions for

the color vector $\hat{n}$ with a given knot topology. We will use in subsequent sections exact analytic knot solutions found in special integrable CP^1 models and generalize them to the case of QCD and effective QCD theory of Skyrme type.

Let us recall first an explicit construction of a simple knot configuration of $\hat{n}$ as a mapping $S^3 \rightarrow S^2$ with unit Hopf number. Surprisingly, such a simple construction leads directly to exact knot solutions found in CP^1 integrable models. Using stereographic projection it is convenient to parameterize the target space S^2 by a complex field $u \in C^1$

$$\hat{n} = \frac{1}{1 + u \bar{u}} \begin{pmatrix} u + \bar{u} \\ -i(u - \bar{u}) \\ u \bar{u} - 1 \end{pmatrix}, \quad (14)$$

A three-dimensional sphere S^3 is given as embedding into R^4 by the following equation

$$|z_1|^2 + |z_2|^2 = 1 \quad (15)$$

where z_1, z_2 are complex coordinates on the complex plane C^2 . The Hopf mapping with Hopf charge $Q_H = 1$ is determined as follows

$$u = \frac{z_1}{z_2}. \quad (16)$$

Starting with a given color vector $\hat{n}$ one can define a magnetic field strength

$$\begin{aligned} H_{\mu\nu} &= \epsilon^{abc} \hat{n}^a \partial_\mu \hat{n}^b \partial_\nu \hat{n}^c = \\ &= \frac{-2i}{(1 + |u|^2)^2} (\partial_\mu u \partial_\nu \bar{u} - \partial_\nu u \partial_\mu \bar{u}). \end{aligned} \quad (17)$$

As we mentioned before, the magnetic field $H_{\mu\nu}$ defines a closed differential two-form, so one can express the dual magnetic Abelian potential (6) in terms of a complex $SU(2)$ doublet $\zeta = (z_1, z_2)$

$$\vec{C}_\mu = -2i\zeta^\dagger \partial_\mu \zeta. \quad (18)$$

For physical implications an explicit realization of Hopf mapping $\hat{n}$ depends on a specified model and topology of four-dimensional space-time. The three-dimensional sphere S^3 in the above definition of Hopf mapping can be related to the physical space-time by several ways. We will consider exact solutions in two cases: the first case, when the sphere S^3 is treated as embedding into Euclidean four-dimensional space-time R^4 (Section III, IV), and the second one, when S^3 is obtained from the real physical space R^3 due to an appropriate compactification by imposing suitable boundary conditions at space infinity for physical fields (Section V, VI). In each case different realizations of the sphere S^3 lead to different topological field configurations in general.

2. Topology of $SU(3)$ QCD vacuum

The consideration of the topological vacuum structure of $SU(2)$ QCD can be generalized to the case of $SU(3)$ gauge group which possesses a more rich topology. The $SU(3)$ gauge invariant Abelian projection is defined as follows [19, 20]

$$\begin{aligned}\vec{A}_\mu &= \hat{A}_\mu + \vec{X}_\mu, \\ \hat{A}_\mu &= A_\mu^r \hat{m}_r + \vec{C}_\mu, \\ C_\mu^a &= -f^{abc} \hat{m}_r^b \partial_\mu \hat{m}_r^c \equiv -(\hat{m}_r \times \partial_\mu \hat{m}_r)^a, \end{aligned} \quad (19)$$

where two color octet fields $\hat{m}_r$, $r = (3, 8)$ correspond to Cartan algebra generators of the Abelian subgroups $U_3(1)$, $U_8(1)$. The group manifold of $SU(3)$ can be described within the formalism of fiber bundle by several ways. One can realize $SU(3)$ manifold as a fiber bundle with the base S^5 and fiber S^3 . This is a generalization of the Hopf fibering $S^1 \rightarrow S^3 \rightarrow S^2$ to the case of $SU(3)$ group. Another possibility is to consider the group manifold $SU(3)$ as a fiber bundle over the coset $M^6 \equiv SU(3)/U(1) \times U(1)$ with a fiber as torus $T^2 = S^1 \times S^1$. The coset space M^6 itself can be treated as a bundle which is locally isomorphic to $CP^2 \times CP^1$. We will consider the last realization of $SU(3)$ manifold because of importance of the Cartan algebra of $SU(3)$ in connection with confinement phenomenon in QCD.

Due to presence of the non-trivial homotopy groups

$$\begin{aligned}\pi_2(SU(3)/U(1) \times U(1)) &= \pi_2(CP^2) \times \pi_2(CP^1) \\ &= Z \times Z, \\ \pi_3(SU(3)/U(1) \times U(1)) &= \pi_3(CP^1) = Z, \end{aligned} \quad (20)$$

one concludes that nonequivalent topological classes of color vector fields $\hat{n}_{3,8}$ are classified by the Hopf number Q_H and two monopole charges (m, n) . As we will see later, the Hopf number is actually determined by two integer winding numbers as well. So that, the classical vacuum topology is described by four independent integers.

Let us construct vacuum configurations in terms of independent complex field variables in analogy with the case of $SU(2)$ QCD. One needs to parameterize the coset M^6 by three complex coordinates. To do this one should express the color vectors $\hat{n}_{3,8}$ in terms of two complex triplet fields Ψ, Φ which are projective coordinates on almost complex homogeneous manifold M^6 . Let us first express the lowest weight vector $\hat{m}_8^a$ introducing a complex triplet field Ψ that parameterizes the coset $CP^2 \simeq SU(3)/SU(2) \times U(1)$ with a maximal stability subgroup

$$\begin{aligned}\hat{m}_8^a &= -\frac{\sqrt{3}}{2} \bar{\Psi} \lambda^a \Psi, \\ \bar{\Psi} \Psi &= 1. \end{aligned} \quad (21)$$

The definition of the vector $\hat{m}_8$ is consistent with the

normalization condition and symmetric d -product operation in the Lie algebra of $SU(3)$

$$\hat{m}_8^2 = 1, \quad d^{abc} \hat{m}_8^b \hat{m}_8^c = -\frac{1}{\sqrt{3}} \hat{m}_8^a. \quad (22)$$

To construct a second Cartan vector $\hat{m}_3$ orthogonal to $\hat{m}_8$ it is convenient to define projectional operators

$$\begin{aligned}P_{\parallel}^{ab} &= \hat{m}_8^a \hat{m}_8^b, \\ P_{\perp}^{ab} &= \delta^{ab} - \hat{m}_8^a \hat{m}_8^b. \end{aligned} \quad (23)$$

With this the vector $\hat{m}_3$ can be parameterized as follows

$$\hat{m}_3^a = P_{\perp}^{ab} \bar{\Phi} \lambda^b \Phi = \bar{\Phi} \lambda^a \Phi + \frac{1}{2} \bar{\Psi} \lambda^a \Psi, \quad (24)$$

where we have introduced a second complex triplet field Φ . The definition of color vectors $\hat{m}_{3,8}$ by Eqs. (21), (24) is invariant under local $\tilde{U}(1) \times \tilde{U}'(1)$ gauge transformations

$$\begin{aligned}\Psi &\rightarrow \exp[i\tilde{\alpha}(x)]\Psi, \\ \Phi &\rightarrow \exp[i\tilde{\alpha}'(x)]\Phi, \end{aligned} \quad (25)$$

which represent explicitly the dual Abelian magnetic symmetry in $SU(3)$ QCD. The dual magnetic potentials $\tilde{C}_\mu^r$ can be expressed through the complex fields

$$\begin{aligned}\tilde{C}_\mu^3 &= 2i(\bar{\Phi} \partial_\mu \Phi + \frac{1}{2} \bar{\Psi} \partial_\mu \Psi), \\ \tilde{C}_\mu^8 &= 2i(-\frac{\sqrt{3}}{2} \bar{\Psi} \partial_\mu \Psi). \end{aligned} \quad (26)$$

One can verify that $\hat{m}_r$ satisfy the following relations

$$\begin{aligned}\hat{m}_r^a \hat{m}_s^a &= \delta_{rs}, \\ d^{abc} \hat{m}_r^b \hat{m}_s^c &= d_{rsq} \hat{m}_q^a, \end{aligned} \quad (27)$$

which imply the orthogonality condition for the complex fields $\bar{\Psi} \Phi = 0$.

Let us define $\Psi = (\psi_1, \psi_2, \psi_3)$ and $\Phi = (\phi_1, \phi_2, \phi_3)$. Introducing projective coordinates

$$\begin{aligned}u_1 &= \frac{\psi_1}{\psi_2}, & u_2 &= \frac{\psi_3}{\psi_2}, \\ v_1 &= \frac{\phi_1}{\phi_2}, & v_2 &= \frac{\phi_3}{\phi_2}, \end{aligned} \quad (28)$$

one can obtain explicit parametrization for the vector $\hat{m}_8$

$$\hat{m}_8 = \frac{1}{1 + |u_1|^2 + |u_2|^2} \cdot \begin{pmatrix} u_1 + u_1^* \\ i(u_1 - u_1^*) \\ |u_1|^2 - 1 \\ u_1^* u_2 + u_2^* u_1 \\ -i(u_1^* u_2 - u_2^* u_1) \\ u_2 + u_2^* \\ -i(u_2 - u_2^*) \\ 1 + |u_1|^2 - 2|u_2|^2 \end{pmatrix}, \quad (29)$$

An explicit expression for the vector $\hat{m}_3$ can be obtained by using Eqn. (24) in a similar manner. Notice, that due to the orthogonality condition $\bar{\Psi}\Phi = 0$ one has an additional constraint

$$1 + {}^*u_1 v_1 + {}^*u_2 v_2 = 0. \quad (30)$$

So that, we can choose three independent complex parameters, for instance, u_1, u_2, v_1 , which contain the whole information on the topology of the space $SU(3)/U(1) \times U(1)$. This can be useful in search of essentially $SU(3)$ solutions with various combinations of topological Hopf and monopole charges corresponding to topologies of the fields u_1, u_2, v_1 .

III. EXACT NON-STATIC SOLUTION IN CP^1 MODEL

In this section we consider exact non-static solution in a simple CP^1 model. The Lagrangian of the CP^1 model with $SU(2)$ triplet field $\hat{n}$ coincides formally with the Lagrangian of the restricted QCD with a vanishing "electric" gauge potential A_μ

$$\mathcal{L} = -\frac{1}{4}H_{\mu\nu}^2(\hat{n}). \quad (31)$$

We use four dimensional spherical coordinate system with Euler angles

$$\begin{aligned} x_1 &= \rho \cos \frac{\theta}{2} \cos \phi, \\ x_2 &= \rho \cos \frac{\theta}{2} \sin \phi, \\ x_3 &= \rho \sin \frac{\theta}{2} \cos \psi, \\ x_4 &= \rho \sin \frac{\theta}{2} \sin \psi, \\ 0 &\leq \theta \leq \pi, \quad 0 \leq \phi, \psi \leq 2\pi. \end{aligned} \quad (32)$$

This coordinate system is suitable especially in studying solutions in Euclidean QCD in the subsequent section.

To define Hopf fibering one introduces complex coordinates z_1, z_2 in the complex plane C^2 (R^4)

$$\begin{aligned} z_1 &= x_1 + ix_2, \\ z_2 &= x_3 - ix_4. \end{aligned} \quad (33)$$

A three-dimensional sphere with the radius ρ is defined by the equation

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 = \rho^2. \quad (34)$$

The Hopf mapping with a unit Hopf charge is defined by

$$u = \frac{z_1}{z_2} = \cot \frac{\theta}{2} e^{i(\phi+\psi)}. \quad (35)$$

Surprisingly, that simple configuration with a Hopf number $Q_H = 1$ represents an exact solution to the equation

of motion for the Lagrangian (31) when the field $\hat{n}$ is defined on three-dimensional sphere S^3 with a fixed radius $\rho = \text{const}$

$$\partial_\mu (\sqrt{g} \partial_\nu u H^{\mu\nu}) = 0. \quad (36)$$

For higher Hopf numbers the solution is provided by the following ansatz

$$u = g(\theta) e^{i(m\psi + n\phi)}. \quad (37)$$

Substituting the ansatz into Euler-Lagrange equations corresponding to the Lagrangian $\mathcal{L}$ one obtains the following solution [33]

$$g(\theta) = \frac{\sqrt{-mn(1+2c_2) + c_1 \log(m^2 + n^2 - 2mn \cos \theta)}}{\sqrt{2mnc_2 - c_1 \log(m^2 + n^2 - 2mn \cos \theta)}}, \quad (38)$$

where the integration constants (c_1, c_2) are fixed by boundary conditions for $\hat{n}$

$$\begin{aligned} \hat{n}(\theta = \pm\pi) &= (0, 0, \pm 1), \\ c_1 &= -\frac{m^2 - n^2}{\log[m^2] - \log[n^2]}, \\ c_2 &= -\frac{\log[2n^2]}{\log[m^2] - \log[n^2]}. \end{aligned} \quad (39)$$

This implies the Hopf charge to be determined by two winding numbers

$$Q_H = mn. \quad (40)$$

In the case $m = n = 1$ the solution reduces to a special one, Eq. (35). Notice, the solution (38) can be treated as a "static" soliton in a sense that it does not depend on the four-dimensional radius ρ . This solution coincides exactly with the solution obtained in [33] with the field $\hat{n}$ defined on three-dimensional sphere of finite radius r_0 . In other words, the topology of the Euclidean space-time is chosen to be $S^3 \times R^1$. The presence of the dimensional parameter r_0 allows to overcome the restrictions of the Derrick theorem to existence of finite energy soliton solutions. Notice, that solution (38) does not admit analytical continuation to the Minkowski space-time.

In our approach we do not restrict the Euclidean four-dimensional space-time to the topology $S^3 \times R^1$. To find a non-static solution to equations of motion of QCD in Euclidean space-time R_E^4 we apply the ansatz used in [33]

$$u = f(\rho, \theta) \exp[i(m\phi + n\psi)]. \quad (41)$$

After proper changing variable

$$g(\rho, \theta) = \frac{1}{1 + f^2(\rho, \theta)} \quad (42)$$

one obtains a linear partial differential equation

$$\rho \partial_\rho (\rho \partial_\rho g) + \frac{4}{A \sin \theta} \partial_\theta (A \sin \theta \partial_\theta g) = 0, \\ A \equiv m^2 \sec^2 \frac{\theta}{2} + n^2 \csc^2 \frac{\theta}{2}. \quad (43)$$

The equation admits separation of variables

$$g(\rho, \theta) = R(\rho)Y(\theta). \quad (44)$$

A solution for the radial function is given by

$$R(\rho) = C_1 \sin[w^2 \log(\mu\rho)] + C_2 \cos[w^2 \log(\mu\rho)], \quad (45)$$

where w^2 is the separation constant. The equation for the angular function $Y(\theta)$

$$\frac{4}{A \sin \theta} \partial_\theta (A \sin \theta \partial_\theta Y(\theta)) - w^2 Y(\theta) = 0 \quad (46)$$

can be reduced to the known Heun equation. Notice, the angular part of our solution is the same as in [33], however, the radial function $R(\rho)$ is different. An essential difference is that our solution has a dimensional mass scale parameter μ which is completely arbitrary contrary to the case of the solution with a fixed radius r_0 in the CP^1 model on the space with topology $S^3 \times R^1$ [33].

IV. KNOT ANSATZ FOR SOLUTIONS IN QCD IN EUCLIDEAN SPACE-TIME

One method of constructing solutions is to perform deformation of vacuum solutions in the theory. Since the vacuum can be described in terms of color vector $\hat{n}$ which has non-trivial topological properties, an interesting possibility arises: to obtain new solutions starting with known solutions with non-trivial Hopf topology of $\hat{n}$ in CP^1 model. Another advantage of using the color field $\hat{n}$ with non-trivial topology in studying vacuum excitations appears when one considers $SU(3)$ QCD where the topological structure becomes more complicate. In this section we will apply the (m, n) -family of the static solutions $\hat{n}$ [33] to construct an ansatz for search of possible new classical solutions in Euclidean $SU(2)$ QCD. We will use four dimensional spherical coordinates with Euler angles. In these coordinates the 't Hooft instanton solution can be reproduced easily from the knot configuration with a unit Hopf charge.

We define a following ansatz with three radial trial functions $f_i(\rho)$

$$\vec{A}_\mu = -f_1(\rho) \tilde{C}_\mu \hat{n} - f_2(\rho) \hat{n} \times \partial_\mu \hat{n} + f_3(\rho) \partial_\mu \hat{n}. \quad (47)$$

Substituting the ansatz into the full equations of motion of QCD

$$\vec{D}_\mu \vec{F}^{\mu\nu} = 0 \quad (48)$$

one can obtain solutions with finite and infinite energy. We select the most interesting solutions:

1. 't Hooft instanton:

$$f_1 = f_2 = \frac{a^2}{a^2 + \rho^2}, \quad f_3 = 0. \quad (49)$$

One can check that the Chern-Simons number for the gauge potential is equalled exactly to the Hopf number $\hat{n}$.

2. There are several gauge equivalent representations for the known 't Hooft instanton. One possible representation is given by

$$f_1 = \frac{a^2}{a^2 + \rho^2}, \quad f_2 = 2 - f_1, \quad f_3 = 0. \quad (50)$$

Notice, the solution represents vacuum excitation over the vacuum configuration given by $f_1 = 0, f_2 = 2$. Such a vacuum corresponds to the vacuum of extended QCD, whereas the vacuum solution with $f_1 = f_2 = 0$ belongs to restricted QCD.

3. Infinite energy solutions

$$f_1 = \frac{1}{2}, \quad f_2 = 1 \pm \frac{1}{2\sqrt{1+c_3}}, \\ f_3 = \pm \frac{c_3}{2\sqrt{1+c_3}}. \quad (51)$$

Radial energy density (density of the Euclidean action) takes the form

$$\mathcal{E}(\rho) = \frac{3 \sin \theta}{2\rho} \quad (52)$$

which is exactly the energy of the knot solution with unit Hopf charge on three-dimensional sphere with radius ρ . In this case it is evident that a solution of the CP^1 model on three dimensional sphere turns into the solution in full Yang-Mills theory.

4. Singular solution with an arbitrary dimensional parameter a

$$f_1 = \frac{a^2}{a^2 + \rho^2}, \quad f_2 = 1, \quad f_3 = 0. \quad (53)$$

The radial energy density is given by

$$\mathcal{E}(\rho) = \frac{16a^4 \sin \theta}{\rho^5}. \quad (54)$$

Notice, the presence of the trial function f_3 leads to increasing the gauge freedom conditioned by gauge transformation which invokes degrees of freedom related to the off-diagonal gluons. As an example, one can find another gauge equivalent instanton solution with a non-vanishing trial function f_3

$$f_1 = \frac{r^2}{a^2 + r^2}, \quad f_2 = 1 \pm \frac{a^2}{\sqrt{2}(a^2 + r^2)}, \\ f_3 = \pm \frac{a^2}{\sqrt{2}(a^2 + r^2)}. \quad (55)$$

To prove the gauge equivalence one can verify that Casimir invariants of the solutions are the same. In the case of $SU(3)$ QCD the adding the third term into the ansatz (47) can be useful in search of a complete set of classical solutions like monopoles which form a non-trivial representation of the Weyl group of $SU(3)$.

V. SINGULAR SOLUTIONS IN QCD IN MINKOWSKI SPACE-TIME

It is known that Wu-Yang monopole represents a solution to classical equations of motion in $SU(2)$ Yang-Mills theory. Within the formalism of the restricted QCD the Wu-Yang monopole is given by the unit color vector directed along the radius

$$\hat{n}^i = \frac{x^i}{r} \quad (i = 1, 2, 3). \quad (56)$$

The solution is singular, it satisfies the equations of motion $\partial^\mu H_{\mu\nu} = 0$ everywhere except the origin. The unit monopole charge of Wu-Yang monopole is provided by the nontrivial homotopy group $\pi_2(SU(2)/U(1))$.

Let us consider a vector $\hat{n}$ with a natural knot topology induced by the definition of the three-dimensional sphere obtained by compactification of the space R^3 to three-sphere S^3 by identifying all points at infinity. To do this we apply stereographic projection of S^3 to R^3

$$\begin{aligned} x_1 &= \frac{2a^2}{a^2 + r^2} X, \\ x_2 &= \frac{2a^2}{a^2 + r^2} Y, \\ x_3 &= \frac{2a^2}{a^2 + r^2} Z, \\ r^2 &\equiv X^2 + Y^2 + Z^2, \end{aligned} \quad (57)$$

where X, Y, Z are Cartesian coordinate in R^3 . The three-sphere S^3 is defined by the equation

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 = a^2, \quad (58)$$

where we keep the radius a of the sphere as a free parameter which will be useful in further. One can pass from the Cartesian coordinate system X, Y, Z to the standard spherical coordinates r, θ, ϕ . The complex coordinates z_1, z_2 can be written as follows

$$\begin{aligned} z_1 &= x_1 + ix_2 = \frac{2a^2 r \sin \theta e^{i\phi}}{a^2 + r^2}, \\ z_2 &= x_3 + ix_4 = \frac{a(2ar \cos \theta - i(a^2 - r^2))}{a^2 + r^2}. \end{aligned} \quad (59)$$

It is convenient to define the Hopf mapping by

$$u = \frac{z_2}{z_1} = \frac{2ar \cos \theta - i(a^2 - r^2)}{2ar \sin \theta} e^{i\phi}. \quad (60)$$

One can write down an explicit expression for the color vector $\hat{n}$ as well

$$\hat{n} = \frac{1}{a^2 + r^2} \cdot \begin{pmatrix} 4ar(2ar \cos \phi \cos \theta + (a^2 - r^2) \sin \phi) \sin \theta \\ 4ar((-a^2 + r^2) \cos \phi + 2ar \sin \phi \cos \theta) \sin \theta \\ (a^2 - r^2)^2 + 4a^2 r^2 \cos(2\theta) \end{pmatrix}. \quad (61)$$

Notice, that magnetic field components have simple expressions and correspond to magnetic helical vortex configuration

$$\begin{aligned} H_{r\theta} &= -\frac{32a^3 r^2 \sin \theta}{(a^2 + r^2)^3}, \\ H_{\theta\phi} &= -\frac{8a^2 r^2 \sin(2\theta)}{(a^2 + r^2)^2}, \\ H_{r\phi} &= -\frac{8a^2 r(a^2 - r^2)(1 - \cos(2\theta))}{(a^2 + r^2)^3}. \end{aligned} \quad (62)$$

Surprisingly, even though the magnetic field is axially symmetric, the energy density $\mathcal{E}$ is spherically symmetric [29], and the knot configuration has a finite energy

$$E = \int d^3V \mathcal{E} = \int r^2 \sin \theta dr d\theta d\phi \frac{256a^4}{(a^2 + r^2)^4} = \frac{32\pi^2}{a}. \quad (63)$$

The solution was first found as an exact solution in the Nicole model with a Lagrangian [29]

$$\mathcal{L} \simeq -((\partial_\mu \hat{n})^2)^{3/2}, \quad (64)$$

and later as a special solution with parameters $m = n = 1$ in the integrable Aratyn-Ferreira-Zimmerman (AFZ) model defined by the following Lagrangian [30]

$$\mathcal{L}_{AFZ} = -\frac{1}{4}(H_{\mu\nu}^2)^{3/4}. \quad (65)$$

We introduce a following axially symmetric ansatz for the restricted gauge potential

$$\begin{aligned} \vec{A}_0^a &= 0, \\ \vec{A}_r^a &= P(r, \theta) \hat{n}^a + \vec{C}_r^a, \\ \vec{A}_\theta^a &= Q(r, \theta) \hat{n}^a + \vec{C}_\theta^a, \\ \vec{A}_\phi^a &= R(r, \theta) \hat{n}^a + \vec{C}_\phi^a, \end{aligned} \quad (66)$$

where P, Q, R are trial functions. Due to the dual $U(1)$ symmetry one can impose a gauge condition on the trial functions. It is suitable to choose a condition $P(r, \theta) = 0$. The ansatz implies the following expression for the scalar

magnetic field $(m, n = r, \theta, \phi)$

$$\begin{aligned}
H_{mn} &= H_{mn}^a \hat{n}^a, \\
H &\equiv dx^m \wedge dx^n H_{mn} = \\
dr \wedge d\theta &\left(\frac{32a^3 r^2 \sin \theta}{(a^2 + r^2)^3} + Q_r \right) + \\
dr \wedge d\phi &\left(\frac{16a^2 r(r^2 - a^2) \sin^2 \theta}{(a^2 + r^2)^3} + R_r \right) + \\
d\theta \wedge d\phi &\left(-\frac{8a^2 r^2 \sin(2\theta)}{(a^2 + r^2)^3} + R_\theta \right). \quad (67)
\end{aligned}$$

After substituting the ansatz into the equations of motion $\vec{D}^\mu \vec{F}_{\mu\nu} = 0$ one can find that all nine equations reduce to three linear partial differential equations

$$\begin{aligned}
(a^2 + r^2)^3 (\sin \theta Q_{r\theta} + \cos \theta Q_r) + 32a^3 r^2 \sin(2\theta) &= 0, \\
(a^2 + r^2)^4 Q_{rr} + 64a^3 r(a^2 - 2r^2) \sin \theta &= 0, \\
(a^2 + r^2)^4 (\sin \theta (r^2 R_{rr} + R_{\theta\theta}) - \cos \theta R_\theta) & \\
- 32(a^2 r^4 (5a^2 + r^2) \sin^3 \theta) &= 0. \quad (68)
\end{aligned}$$

The fact that we have finally linear differential equations is caused by Abelian structure of the chosen ansatz. The solution to the first two equations is given as follows

$$\begin{aligned}
Q &= c_1 + c_2 \frac{r}{\sin \theta} - \frac{4ar(a^2 - r^2)}{(a^2 + r^2)^2} \sin \theta \\
&- 4 \arctan\left(\frac{r}{a}\right) \sin \theta. \quad (69)
\end{aligned}$$

A solution to the equation for the function R can be found as a sum of a general solution $R_0(r, \theta)$ to the homogeneous part of the equation and a special solution $R_1(r, \theta)$ to the inhomogeneous equation

$$\begin{aligned}
R(r, \theta) &= R_0(r, \theta) + R_1(r, \theta), \\
R_1(r, \theta) &= \left(\frac{c_{01}}{r} + c_{02} r^2 + \frac{8a^2 r^2}{(a^2 + r^2)^2} \right) \sin^2 \theta. \quad (70)
\end{aligned}$$

The homogeneous equation for $R_0(r, \theta)$ is separable and the general solution is given by

$$\begin{aligned}
R_0(r, \theta) &= \sum_w K(r; w) Y(\theta; w), \\
K(r; w) &= a_w r^{(1-\sqrt{1+4w})/2} + b_w r^{(1+\sqrt{1+4w})/2}, \\
Y(\theta; w) &= c_w {}_2F_1[-p_+, -p_-, 1/2; \cos^2 \theta] + \\
&d_w \cos \theta {}_2F_1[p_-, p_+, 3/2; \cos^2 \theta], \\
p_\pm &= (1 \pm \sqrt{1+4w})/4, \quad (71)
\end{aligned}$$

where w is a constant of separation. The hypergeometric functions ${}_2F_1(a, b, c; z)$ in the above equations reduce to trigonometric polynomials for integer and semi-integer values of $p_\pm$, i.e., if we impose the following conditions

$$\begin{aligned}
w &= l(l+1), \quad l = 0, 1, 2, \dots, \\
c_l &= 0, \quad \text{for } l = 2, 4, 6, \dots, \\
d_l &= 0, \quad \text{for } l = 1, 3, 5, \dots. \quad (72)
\end{aligned}$$

In general, the solution represents a class of singular infinite energy field configurations due to singularities at the origin, $r = 0$, or at space infinity, $r = \infty$. Let us consider special solutions which have singularities at the origin and the energy density $E(r)$ falling down at infinity as $\left(\frac{1}{r}\right)^{\alpha>1}$ ($l = 0$):

(i) $c_1 = c_2 = c_{01} = c_{02} = b_w = 0$, $a_w = d_w = 1$: the solution represents the Wu-Yang monopole.

(ii) $c_1 = c_2 = c_{02} = b_w = 0$, $a_w = d_w = 1$: a singular monopole solution with two non-zero magnetic components $H_{\theta\phi}, H_{r\phi}$.

(iii) if $c_2 \neq 0$: all components of the magnetic field H_{mn} are not vanishing. In this case one has an additional singularity along OZ axis.

VI. SKYRME TYPE MODELS WITH KNOT SOLITONS

The knot soliton (60) represents an exact solution with unit Hopf charge in integrable models [29, 30] with Lagrangians (64, 65) containing fractional degrees of the kinetic terms. Another interesting model was found in [34] where a generalized Faddeev-Skyrme model was considered as a possible low energy effective theory of QCD. In particular, it had been found that the Lagrangian

$$\mathcal{L} = -\frac{1}{4} H_{\mu\nu}^2 + \xi (\partial_\mu \hat{n})^4 \quad (73)$$

leads to existence of analytic solutions with zero energy in the critical case $\xi = 1$. These models are far from physical theories like standard or effective low energy QCD. An interesting question arises, whether a Skyrme type model of effective low energy QCD exists which admits a finite energy exact solution with the knot configuration (60). It turns out that there is a family of Skyrme type models with an additional scalar field which admits exact analytic solutions for different Hopf numbers.

Let us consider the following Lagrangian

$$\mathcal{L}(\phi, \hat{n}) = -\mu^2 (\partial_\mu \phi)^2 - \frac{\beta}{4} \phi^2 (\partial_\mu \hat{n})^2 - \frac{\nu}{32} \frac{1}{\phi^2} H_{\mu\nu}^2. \quad (74)$$

The term proportional to $\frac{1}{\phi^2} (\partial_\mu \hat{n})^4$ can be added to the Lagrangian as well. We will not consider this case since the structure of the exact solution is not affected in principal by adding such a term. Up to change of variables the first two terms in (74) coincide with the respective first two terms in the original Skyrme model. The scalar field ϕ can be interpreted as a fluctuation of the length of the color field $\hat{n}$.

The exact solution in AFZ model (65) in the case $m = n$ in toroidal coordinates has the following form [30]

$$u(\eta, \xi, \phi) = \frac{1}{\sinh \eta} \exp[im(\xi + \phi)]. \quad (75)$$

The Hopf charge of the solution is given by $Q_H = m^2$. The solution with Hopf charge $Q_H = 1$ coincides with the knot configuration (60) in spherical coordinates. We have found for the case $m = n = 1, \dots, 5$ that for each topological charge Q_H there is a special Lagrangian (74) with constrained parameters μ^2, β, ν which admits an exact finite energy solution. The solution for the scalar field ϕ is the same for various Q_H

$$\phi = \frac{a}{\sqrt{a^2 + r^2}}, \quad (76)$$

and the solution for $\hat{n}$ is given by Eq. (60). The constraints on the parameters μ^2, β, ν for topological charges $Q_H = m^2$ are given as follows:

$Q_H = 1$:

$$\frac{3}{8}\mu^2 + \beta - \frac{\nu}{a^2} = 0. \quad (77)$$

In the case of $\nu = 0$ one has a model without any dimensional parameters while possessing a static solution which includes a dimensional parameter a corresponding to the size of the knot. This solution represents a saddle point type solution with spherically symmetric radial energy density

$$\mathcal{E}_0 = \frac{a^2\mu^2(r^2 - 3a^2)}{(a^2 + r^2)^3}. \quad (78)$$

One can verify that the total energy after integration over the space vanishes identically. The solution is rather unphysical since the Hamiltonian is not positively defined. Let us choose another condition on the parameters, $\beta = 0$. In this case the energy density becomes positively defined

$$\mathcal{E}_1 = \frac{8\nu r^2}{3(a^2 + r^2)^3}. \quad (79)$$

This implies a finite total energy

$$E_1 = \frac{2\pi^2}{a}. \quad (80)$$

For Hopf charges $Q_H > 1$ corresponding to winding numbers $m = n = 2, 3, 4, 5$ one can find respective exact solutions under the following condition

$$\beta = 0, \quad \mu^2 - \frac{8m^2\nu}{3a^2} = 0. \quad (81)$$

The radial energy density is

$$\mathcal{E}_m = \frac{8m^2\nu(r^2 + 3a^2)}{3(a^2 + r^2)^3}. \quad (82)$$

The total energy of the solution corresponding to each model with a Lagrangian specified by parameters β, μ, ν is proportional to the Hopf charge

$$E_m = \frac{4\pi^2\nu}{a}Q_H. \quad (83)$$

For the case $m \neq n$ we expect that knot solutions in such models will imply the total energy expressed by the same equation (83).

VII. DISCUSSION

In standard approach to description of the classical vacuum in QCD in terms of pure gauge connection all topologically non-equivalent vacuums are classified by Chern-Simons number. Using an explicit construction of the classical vacuum in terms of the color vector $\hat{n}$ we have demonstrated that classical vacuum has a more rich topological structure. Namely, all non-trivial topological vacuums are described by homotopy groups $\pi_{2,3}(SU(N)/H)$, with H being a maximal Abelian subgroup of $SU(N)$. In the present paper we concern mainly with knot topology of the classical vacuum in the case of $SU(2)$ QCD. A natural question arises on possible physical manifestations of such topological vacuum structure. We have considered excitations over the vacuum with non-trivial Hopf charge to find new analytic classical solutions in the CP^1 model, standard QCD and Skyrme type low energy effective theory of QCD. Our method of studying new solutions maybe useful in search of essentially $SU(3)$ topological solitons like instantons and monopoles. It is interesting to study possible physical applications of the proposed Skyrme type model which can be relevant to the low energy effective theory of QCD like the Faddeev-Skyrme model [11].

Certainly, since QCD is essentially a quantum theory, it is of great importance to find physical effects of the non-trivial topological structure at quantum level where the degeneracy of the classical vacuum should disappear. One possible effect includes a non-trivial Weyl symmetric structure of the minimums in two-loop effective potential corresponding to presence of monopole condensates [28]. Another implication of the knot field configurations in QCD is to study effects related with contributions of all possible knot topologies to physical quantities like the Wilson loop functional and effective action. In Euclidean sector the contribution of non-static solutions may play an important role as well.

Acknowledgments

Authors are grateful to Prof. Y. M. Cho for numerous discussions. One of authors (D.G.P.) thanks E. N. Tsoy and R. M. Galimzyanov for useful discussions. The work is supported by NSFC (Grants 11035006 and 11175215), CAS (Contract No. 2011T1J31), and by UzFFR (Grant F2-FA-F116).

-
- [1] T. Schafer, E. V. Shuryak, *Rev. Mod. Phys.* **70**, 323 (1998).
 - [2] Y. Nambu, *Phys. Rev.* **D10**, 4262 (1974).
 - [3] S. Mandelstam, *Phys. Rep.* **23C**, 245 (1976).
 - [4] A. Polyakov, *Nucl. Phys.* **B120**, 429 (1977).
 - [5] G. K. Savvidy, *Phys. Lett.* **B71**, 133 (1977).
 - [6] N. K. Nielsen and P. Olesen, *Nucl. Phys.* **B144**, 485 (1978).
 - [7] Y. M. Cho and D. G. Pak, *Phys. Rev.* **D65**, 074027 (2002).
 - [8] P. Paccioli and E. Shuryak, QCD Topology at Finite Temperature: Statistical Mechanics of Selfdual Dyons, arXiv: 1301.2523 [hep-ph].
 - [9] T. H. R. Skyrme, *Proc. Roy. Soc. Lond.* **A260**, 127 (1961).
 - [10] N. Manton and P. Sutcliffe, *Topological solitons*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, (2004).
 - [11] L. D. Faddeev and A. J. Niemi, *Nature* **387**, 58 (1997).
 - [12] Y. M. Cho, *Phys. Rev. Lett.* **87**, 252001 (2001).
 - [13] A. Belavin, A. Polykov, A. Schwartz, and Y. Tyupkin, *Phys. Lett.* **59B**, 85 (1975).
 - [14] C. Callan, R. Dashen, and D. Gross, *Phys. Lett.* **B63**, 334 (1976).
 - [15] R. Jackiw and C. Rebbi, *Phys. Rev. Lett.* **37**, 172 (1976).
 - [16] P. Goddard, J. Nuyts and D. Olive, *Nucl. Phys.* **B125**, 1 (1977).
 - [17] Y. M. Cho, *Phys. Rev. Lett.* **44**, 1115 (1980).
 - [18] Y. M. Cho, *Phys. Lett.* **B115**, 125 (1982).
 - [19] Y. M. Cho, *Phys. Rev.* **D21**, 1080 (1980).
 - [20] Y. M. Cho, *Phys. Rev. Lett.* **46**, 302 (1981).
 - [21] Y. S. Duan and Mo-Lin Ge, *Sci. Sinica* **11**, 1072 (1979).
 - [22] Y.-S. Duan, P. M. Zhang, *Mod. Phys. Lett.* **A17**, 2283 (2002).
 - [23] L. D. Faddeev and A. J. Niemi, *Phys. Rev. Lett.* **82** 1624 (1999).
 - [24] L. D. Faddeev, A. J. Niemi, *Phys. Lett.* **B449**, 214 (1999).
 - [25] S. V. Shabanov, *Phys. Lett.* **B458**, 322 (1999).
 - [26] P. van Baal and A. Wipf, *Phys. Lett.* **B515**, 181 (2001).
 - [27] Y. M. Cho, *Phys. Lett.* **B644**, 208 (2007).
 - [28] Y. M. Cho, D. G. Pak, P. M. Zhang, L. P. Zhou, *Phys. Rev.* **D86** 045025 (2012).
 - [29] D. A. Nicole, *J. Phys.* **G4**, 1363 (1978).
 - [30] H. Aratyn, L. A. Ferreira and A. H. Zimerman, *Phys. Rev. Lett.* **83**, 1723 (1999).
 - [31] J. Evslin, S. Giacomelli, *JHEP* 1104:022 (2011).
 - [32] D. G. Pak, Y. M. Kim and T. Tsukioka, *Phys. Rev.* **D85**, 084006 (2012).
 - [33] E. De Carli, L. A. Ferreira, *J. Math. Phys.* **46** 012703 (2005).
 - [34] L. A. Ferreira, A. C. Riserio do Bonfim, *JHEP* 1003:119 (2010).